

# Non-commutative geometry and covariance: from the quantum plane to quantum tensors\*

J.A. de Azcárraga<sup>†</sup>, P.P. Kulish<sup>1†</sup> and F. Rodenas<sup>†‡</sup>

<sup>†</sup> *Departamento de Física Teórica and IFIC,  
Centro Mixto Universidad de Valencia-CSIC  
E-46100-Burjassot (Valencia), Spain.*

<sup>‡</sup> *Departamento de Matemática Aplicada,  
Universidad Politécnica de Valencia  
E-46071 Valencia, Spain.*

## Abstract

Reflection and braid equations for rank two  $q$ -tensors are derived from the covariance properties of quantum vectors by using the  $R$ -matrix formalism.

## 1 Introduction

Quantum groups may be looked at in various ways. From a mathematical point of view, they may be introduced by making emphasis on their  $q$ -deformed enveloping algebra aspects [1, 2] or by making emphasis in the  $R$ -matrix formalism that describe the deformed group algebra [3]. A point of view which is particularly useful in possible physical applications is to look at quantum groups as symmetries of *quantum spaces* [4, 5]. The simplest example of this approach is constituted by the well known quantum plane  $C_q^2$ , or associative *algebra* (a  $q$ -plane is not a manifold) generated by two elements  $(x, y) = X$  (a  $q$ -two-vector) subjected to the commutation property [4]

$$xy = qyx \quad . \quad (1)$$

---

<sup>1</sup>On leave of absence from the St.Petersburg's Branch of the Steklov Mathematical Institute of the Russian Academy of Sciences.

\*Lecture delivered (by J.A.) at the 3rd Colloquium 'Quantum groups and Physics', Prague, June 1994. To appear in the proceedings (Czechoslovak J. Phys.).

The commutation relation (1) can also be expressed by using the  $q$ -symplectic metric  $\epsilon^q$

$$\epsilon^q = \begin{pmatrix} 0 & q^{-1/2} \\ -q^{1/2} & 0 \end{pmatrix} , \quad (\epsilon^q)^2 = -I \quad (2)$$

by the equation

$$X^t \epsilon^q X = 0 \quad , \quad \epsilon_{ij}^q X_i X_j = 0 \quad (3)$$

which reflects that the  $q$ -symplectic norm of a  $q$ -two-vector vanishes.

It is also possible to introduce a pair of (odd) variables  $(\xi, \eta) = \Omega$  (an odd  $q$ -two-vector) satisfying

$$\xi \eta = -\frac{1}{q} \eta \xi \quad , \quad \xi^2 = 0 = \eta^2 \quad . \quad (4)$$

If it is required that, after the transformations  $X' = TX$ ,  $\Omega' = T\Omega$  the new entities  $(x', y')$ ,  $(\xi', \eta')$  satisfy also (1), (4), it is found that the commutation properties of the elements of  $T$

$$T = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad (5)$$

are completely determined. These are the well known relations (the entries of  $T$  commute with those of  $X$  and  $\Omega$ )

$$\begin{aligned} ab &= qba \quad , & ac &= qca \quad , & bd &= qdb \quad , \\ cd &= qdc \quad , & [a, d] &= \lambda cb \quad , & [b, c] &= 0 \quad , \end{aligned} \quad (6)$$

( $\lambda \equiv q - q^{-1}$ ) which constitute a presentation of the  $GL_q(2, C)$  algebra generated by  $(a, b, c, d)$ . For  $q=1$ ,  $(x, y)$  commute and  $(\xi, \eta)$  anticommute. In a non-commutative differential calculus this second set of variables are identified with the differentials ([5]) of  $(x, y)$ . Here we shall consider the quantum plane (1) only as the representation (co-module) space of the  $GL_q(2, C)$  quantum group (6). In terms of the  $R$ -matrix formalism [3], eqs. (1) and (6) may be written as (see eq. (11) below)

$$R_{12} X_1 X_2 = q X_2 X_1 \quad \Longleftrightarrow \quad R_{21}^{-1} X_1 X_2 = q^{-1} X_2 X_1 \quad , \quad (7)$$

$$R_{12} T_1 T_2 = T_2 T_1 R_{12} \quad \Longleftrightarrow \quad R_{21}^{-1} T_1 T_2 = T_2 T_1 R_{21}^{-1} \quad , \quad (8)$$

where the standard notation  $T_1 = T \otimes \mathbf{1}$ ,  $T_2 = \mathbf{1} \otimes T$  ( $T_{1ij,kl} = T_{ik} \delta_{jl}$ ,  $T_{2ij,kl} = \delta_{ik} T_{jl}$ ,  $i, j, k, l = 1, 2$ ) has been used, and  $X_1 X_2$  and  $X_2 X_1$  are, respectively, the four-vectors  $(xx, xy, yx, yy)$  and  $(xx, yx, xy, yy)$ . Both relations in (7) (and in (8)) are equivalent. This is easy to see by using the permutation operator  $\mathcal{P}$  which gives  $(\mathcal{P} R \mathcal{P})_{ij,kl} = R_{ji,lk}$  ( $\mathcal{P} R_{12} \mathcal{P} = R_{21}$ ) and  $(\mathcal{P} X_1 X_2)_{ij} = (X_1 X_2)_{ji}$  *i.e.*,  $\mathcal{P} X_1 X_2 = X_2 X_1$ . In this matrix notation it is obvious that (8) is consistent with the requirement of invariance of (7) under the transformation  $X' = TX$ ,

$R_{12}X'_1X'_2 = qX'_2X'_1$ . Since the elements of  $X$  commute with the entries of  $T$ , we obtain

$$\begin{aligned} R_{12}X'_1X'_2 &= R_{12}(T_1X_1)(T_2X_2) = R_{12}T_1T_2X_1X_2 \\ &= T_2T_1R_{12}X_1X_2 = qT_2T_1X_2X_1 = qX'_2X'_1 \end{aligned} \quad (9)$$

using (8), and the invariance of (7) follows: the preservation of (7) under the ‘ $q$ -symmetry’ transformation requires (8). In components, (7) reads

$$R_{ij,kl}X_kX_l = qX_jX_i \quad , \quad \hat{R}_{ij,kl}X_kX_l = qX_iX_j \quad , \quad (10)$$

where  $R_{12}$  and  $\mathcal{P}$  ( $\hat{R} = \mathcal{P}R$ ,  $\hat{R}_{ij,kl} = R_{ji,kl}$ ) are given by

$$R = \begin{bmatrix} q & & & \\ & 1 & 0 & \\ & \lambda & 1 & \\ & & & q \end{bmatrix} \quad , \quad \mathcal{P} = \begin{bmatrix} 1 & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & 1 \end{bmatrix} \quad . \quad (11)$$

Although the indices in all previous expressions take the values 1, 2, the  $R$ -matrix form of the basic expressions (7) and (8) makes it clear how to generalize them to  $GL_q(n, C)$ ; all that is needed is the appropriate  $n^2 \times n^2$   $R$ -matrix, which is given by [3]

$$R_{ij,kl} = \delta_{ik}\delta_{jl}(1 + \delta_{ij}(q - 1)) + \lambda\delta_{il}\delta_{jk}\theta(i - j) \quad i, j, \dots = 1 \dots n \quad (12)$$

$$\theta(i - j) = \begin{cases} 0 & i \leq j \\ 1 & i > j \end{cases} \quad .$$

With it, the relations defining the ‘quantum hyperplane’

$$X = (x_1, \dots, x_n) \quad , \quad x_i x_j = q x_j x_i \quad (i < j) \quad i, j = 1 \dots n \quad (13)$$

are again expressed by (7) and preserved under  $GL_q(n, C)$  because of (8).

All this is, of course, well known. In this report we exhibit how to extend these  $q$ -vector constructions to higher rank quantum tensors (see also [6, 7]). In particular, we shall consider the simplest example of  $q$ -twistors constructed from  $q$ -two-vectors (spinors) (1), (3), (7) and the application to  $q$ -Minkowski space algebras [8].

## 2 Other covariant objects. Quantum twistors

Consider two isomorphic objects  $X$  and  $Z$ , and their hermitian conjugates  $X^\dagger$  and  $Z^\dagger$ , transforming under the coaction of two different quantum groups  $T$  and  $T^\dagger$  by

$$\begin{aligned} X' &= TX & , & & X'^\dagger &= X^\dagger T^\dagger \\ Z' &= TZ & , & & Z'^\dagger &= Z^\dagger T^\dagger \end{aligned} \quad (14)$$

For instance, in the classical  $SL(2, C)$  case there are two fundamental representations,  $D^{\frac{1}{2}, 0}$  and  $D^{0, \frac{1}{2}}$ , realized by complex unimodular matrices  $A$  and

$(A^{-1})^\dagger$ . In the quantum case this corresponds to taking two copies  $T$  and  $\tilde{T}$  of  $SL_q(2, C)$ , with the obvious ‘reality’ condition added,  $\tilde{T}^{-1} = T^\dagger$ . In the  $q$ -case one has to add the commutation relations between elements of  $T$  and  $T^\dagger$ .

The commutation relations in the general situation involve four  $R$ -matrices  $R^{(i)}$ ,  $i = 1, \dots, 4$ ,

$$\begin{aligned} R^{(1)}T_1T_2 &= T_2T_1R^{(1)} \quad , \\ T_1^\dagger R^{(2)}T_2 &= T_2R^{(2)}T_1^\dagger \quad , \\ T_2^\dagger R^{(3)}T_1 &= T_1R^{(3)}T_2^\dagger \quad , \\ R^{(4)}T_1^\dagger T_2^\dagger &= T_2^\dagger T_1^\dagger R^{(4)} \quad . \end{aligned} \tag{15}$$

The consistency of these equations requires  $R^{(2)} = \mathcal{P}R^{(3)}\mathcal{P} = R^{(3)\dagger}$  and  $R^{(4)} = R^{(1)\dagger}$  or  $R^{(4)} = (\mathcal{P}R^{(1)-1}\mathcal{P})^\dagger$ . Notice that  $R^{(1)}$  and  $R^{(4)}$  are the  $R$ -matrices of two quantum groups  $T$  and  $T^\dagger$  related by a  $*$ -operation (and that  $R^{(1)}$ , *e.g.*, may be taken as  $R_{12}$  or  $R_{21}^{-1}$ ). In contrast,  $R^{(2)}$  (and hence  $R^{(3)}$ ) is a matrix defining how the elements of both quantum groups commute and accordingly it is not a priori fixed. In general, one could introduce instead of  $T^\dagger$  another matrix  $S$ ; the  $q$ -matrices  $S$  and  $T$  need not even have the same dimension. If, say,  $T$  and  $S$  are  $n \times n$  and  $m \times m$  matrices,  $S_1$  and  $T_2$  in the second equation of (15) would be  $S_1^\dagger = (S^\dagger \otimes \mathbf{1}_n)$  and  $T_2 = (\mathbf{1}_m \otimes T)$  and  $R^{(2)}$  would be an  $(m \times n) \times (m \times n)$  matrix. Similarly, in the third equation  $S_2^\dagger = (\mathbf{1}_n \otimes S^\dagger)$ ,  $T_1 = (T \otimes \mathbf{1}_m)$  and  $R^{(3)}$  would be an  $(n \times m) \times (n \times m)$  matrix, while  $R^{(4)}$  would be an  $m^2 \times m^2$  matrix.

The form of the eqs. (15) is the result of the equations which express the commutation relations among the components of the vectors  $X, Z$ . Since in principle  $T$  and  $T^\dagger$  do not commute, we have to allow for possibly non-trivial commutation relations among the components of  $X$  and  $Z^\dagger$ . Thus, the set of commutation relations left invariant is given by

$$\begin{aligned} R^{(1)}X_1X_2 &= \kappa_1X_2X_1 \quad , \\ Z_1^\dagger R^{(2)}X_2 &= X_2Z_1^\dagger \quad , \\ Z_2^\dagger R^{(3)}X_1 &= X_1Z_2^\dagger \quad , \\ \kappa_2Z_1^\dagger Z_2^\dagger &= Z_2^\dagger Z_1^\dagger R^{(4)} \quad , \end{aligned} \tag{16}$$

where  $\kappa_1$  and  $\kappa_2$  are appropriate eigenvalues of the  $R$ -matrices. The invariance of the first and last equations is proven as in Sec.1 and the others similarly. For instance, for the second equation we check that

$$Z_1'^\dagger R^{(2)}X_2' = (Z_1^\dagger T_1^\dagger)R^{(2)}(T_2X_2) = Z_1^\dagger T_2R^{(2)}T_1^\dagger X_2$$

$$= T_2 Z_1^\dagger R^{(2)} X_2 T_1^\dagger = T_2 X_2 Z_1^\dagger T_1^\dagger = X_2' Z_1'^\dagger \quad (17)$$

using the second equations in (15) and (16), respectively, in the second and fourth equalities. In particular, if  $R^{(2)} = I = R^{(3)}$ , both quantum groups are independent (commuting), and this is reflected in the fact that the components of  $X$  and  $Z^\dagger$  commute.

Let us use the above construction to introduce another covariant object which generalizes (with some restrictions) the concept of twistor to the  $q$ -deformed case. Let  $X$  and  $Z^\dagger$  satisfy the previous set of commutation relations. In particular,  $X$  and  $Z^\dagger$  may be, for instance,  $q$ -two-vectors ( $q$ -spinors), of  $SL_q(2, C)$ ; this case will be analyzed in more detail below. Tensoring two  $q$ -vectors we introduce the object

$$K \equiv X Z^\dagger \quad (K_{ij} = X_i Z_j^\dagger) \quad . \quad (18)$$

Then, the transformation of  $K$  induced by (14) is

$$\varphi : K \longmapsto K' = T K T^\dagger \quad (K'_{ij} = T_{im} K_{mn} T_{nj}^\dagger) \quad . \quad (19)$$

The entries of  $K$  are, of course, non-commuting. We shall see that these commutation relations can be expressed in a closed, elegant and compact equation which permits to extract the algebra generated by the entries of  $K$  without considering its explicit realization in terms of the components of  $X$  and  $Z^\dagger$ . Using the above relations we may now derive the equation describing the commutation relations which define the algebra generated by the entries of  $K$ . With  $K_1 = X_1 Z_1^\dagger$  ( $K_{1\,ij,kl} = (K \otimes \mathbf{1})_{ij,kl} = X_i Z_k^\dagger \delta_{jl}$ ) and  $K_2 = X_2 Z_2^\dagger$  ( $K_{2\,ij,kl} = (\mathbf{1} \otimes K)_{ij,kl} = \delta_{ik} X_j Z_l^\dagger$ ), we find using (16) that

$$\begin{aligned} R^{(1)} K_1 R^{(2)} K_2 &= R^{(1)} X_1 Z_1^\dagger R^{(2)} X_2 Z_2^\dagger = R^{(1)} X_1 X_2 Z_1^\dagger Z_2^\dagger \\ &= (\kappa_1/\kappa_2) X_2 X_1 Z_2^\dagger Z_1^\dagger R^{(4)} = (\kappa_1/\kappa_2) X_2 Z_2^\dagger R^{(3)} X_1 Z_1^\dagger R^{(4)} \quad . \end{aligned} \quad (20)$$

Hence, the commuting properties of the quantum twistor are given by

$$R^{(1)} K_1 R^{(2)} K_2 = (\kappa_1/\kappa_2) K_2 R^{(3)} K_1 R^{(4)} \quad (21)$$

Eq. (21) is (with  $\kappa_1/\kappa_2 = 1$ ) nothing else than the reflection equation with no spectral parameter dependence (see [6, 7] and references therein and [9] in the context of braided algebras) which follows by imposing the invariance of the commuting properties of the entries of  $K$  by the coaction (19). As shown here, eq. (21) also follows from interpreting  $K$  as an object made out of two  $q$ -‘vectors’, in general not necessarily of the same dimension so that in general  $K$  is not a squared matrix.

Let  $X, Z$  be two  $q$ -two-vectors (spinors). Then,  $K = X X^\dagger$  is a (null) *quantum twistor*: as we shall see, its quantum determinant ( $\det_q K$ ) is necessarily zero (as it is as well for  $X Z^\dagger$ ). In contrast, the  $q$ -twistor  $K = X Z^\dagger + Z X^\dagger$  has  $\det_q K \neq 0$ .

Notice that, in general, there are four possibilities to write (21) (obviously, related in between) since there are two possibilities for  $R^{(1)}$  and for  $R^{(4)}$  in (15) and in (16) (see (7) and (8)). However, this freedom is reduced when covariant objects  $K$  constructed out of four vectors are considered since covariance requires to introduce commutation relations between  $Z$  and  $X$  and between  $Z^\dagger$  and  $X^\dagger$  using  $R^{(1)}$  and  $R^{(4)}$ . Let us consider the hermitian matrix

$$K = XZ^\dagger + ZX^\dagger \quad (22)$$

( $Z$  and  $X$  have the same number of components). To compute the commutation properties of  $K$ , the complete set of relations among  $X$ ,  $Z$ ,  $X^\dagger$  and  $Z^\dagger$  are required. Thus, besides (16), we need to introduce the following set of covariant relations

$$\begin{aligned} R^{(1)}X_1Z_2 &= Z_2X_1 \quad , \\ Z_1^\dagger R^{(2)}Z_2 &= Z_2Z_1^\dagger \quad , \\ X_2^\dagger R^{(3)}X_1 &= X_1X_2^\dagger \quad , \\ X_1^\dagger Z_2^\dagger &= Z_2^\dagger X_1^\dagger R^{(4)} \quad , \end{aligned} \quad (23)$$

the structure of which is again dictated from (15) by covariance. From the first and the last eqs. in (23) we obtain (supposing  $R^{(1)}$  real)

$$R^{(4)} = (R^{(1)})^t \quad (24)$$

which implies that the eigenvalues are equal,  $\kappa_1 = \kappa_2$ . Then, the commutation relation for the entries of  $K$  (in matrix form) are easily computed using (16) and (23)

$$\begin{aligned} R^{(1)}K_1R^{(2)}K_2 &= R^{(1)}(X_1Z_1^\dagger + Z_1X_1^\dagger)R^{(2)}(X_2Z_2^\dagger + Z_2X_2^\dagger) \\ &= (X_2Z_2^\dagger + Z_2X_2^\dagger)R^{(3)}(X_1Z_1^\dagger + Z_1X_1^\dagger)R^{(4)} \\ &= K_2R^{(3)}K_1R^{(4)} \end{aligned} \quad (25)$$

where  $R^{(1)} = R^{(4)t} = \mathcal{P}R^{(4)}\mathcal{P}$  and  $R^{(2)} = \mathcal{P}R^{(3)}\mathcal{P}$  and Hecke's condition for the  $R$ -matrix has been used. Now, we have only one reflection equation for  $K$ , since the two possibilities for  $R^{(1)}$  produce two equations which are identical after a similarity transformation with  $\mathcal{P}$ .

Notice that  $K$  in (22) is constructed from two parts, each one of them satisfying the same algebra relations (25):

$$K = K^{(1)} + K^{(2)} \quad , \quad K^{(1)} = XZ^\dagger \quad , \quad K^{(2)} = ZX^\dagger \quad . \quad (26)$$

These two pieces have specific commutation properties among themselves. Indeed, the (mixed) commutation relations (23) lead to the following non-commuting property between the matrices  $K^{(1)}$  and  $K^{(2)}$  (non-symmetric under the interchange of  $K^{(1)}$  and  $K^{(2)}$ )

$$R^{(1)}K_1^{(1)}R^{(2)}K_2^{(2)} = K_2^{(2)}R^{(3)}K_1^{(1)}(\mathcal{P}R^{(4)}\mathcal{P})^{-1} \quad . \quad (27)$$

Here  $R^{(4)} = R^{(1)^\dagger}$  and the two possibilities for  $R^{(1)}$  produce two different equations for  $K^{(1)}$  and  $K^{(2)}$  which transform into each other by exchanging  $(1) \leftrightarrow (2)$  in  $K^{(i)}$ . Both had to be possible since  $K = K^{(1)} + K^{(2)}$  is symmetric under this exchange. Equation (27), here obtained from the commutation relations (23), is known as ‘braiding equation’ [9]. The commutation properties among the elements of  $K^{(1)}$  and  $K^{(2)}$  are such that the sum of two objects satisfying (25) verifies also the same relation. Within this terminology, the ‘mixed’ eqs. (23) are the braiding relations for  $q$ -vectors.

From now on, we shall restrict ourselves to the two-dimensional case which will be useful in the application to  $q$ -Minkowski space [8]. We shall start by discussing the

### **$q$ -determinant of $K$ :**

Let the quantum group matrices  $T$  and  $T^\dagger$  be  $2 \times 2$  matrices. There exists an invariant quadratic element from  $K$ , the  $q$ -determinant of  $K$ . It is defined by [6]

$$\det_q K P_- \equiv P_- K_1 \hat{R}^{(3)} K_1 P_- \quad (28)$$

where  $\hat{R}^{(3)} = \mathcal{P}R^{(3)}$  and  $P_-$  is the  $q$ -antisymmetrizer of the  $R$ -matrix corresponding to the quantum groups  $T$  and  $T^\dagger$ . The  $q$ -determinant of  $T$  and  $T^\dagger$  are given by [3]

$$\det_q T P_- = P_- T_1 T_2 \quad , \quad \det_q T^\dagger P_- = T_2^\dagger T_1^\dagger P_- \quad (29)$$

and the projector  $P_-$  can be expressed in terms of the  $q$ -epsilon tensor (2)

$$P_{-ij,kl} = [2]^{-1} \epsilon_{ij}^q \epsilon_{kl}^q \quad , \quad P_- = \frac{1}{[2]} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & q^{-1} & -1 & 0 \\ 0 & -1 & q & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} . \quad (30)$$

When  $(\det_q T)(\det_q T^\dagger) = 1$ ,  $\det_q K$  is invariant under the coaction (19). Using the third eq. in (15) and (29)

$$\begin{aligned} \det_q (TKT^\dagger) &= P_- (T_1 K_1 T_1^\dagger) \hat{R}^{(3)} (T_1 K_1 T_1^\dagger) P_- \\ &= P_- T_1 T_2 K_1 \hat{R}^{(3)} K_1 T_2^\dagger T_1^\dagger P_- \\ &= (\det_q T) (\det_q K) (\det_q T^\dagger) . \end{aligned} \quad (31)$$

Thus, if  $(\det_q T)(\det_q T^\dagger) = 1$  we obtain that  $\det_q (TKT^\dagger) = \det_q K$ . The centrality of  $\det_q K$  requires some YBE-like conditions on the  $R^{(i)}$  ( $i = 1, 2, 3, 4$ ) matrices in (21).

Using the definition (28) and the  $R$ -matrix property  $\hat{R}_{ab,cd}^{(3)} = R_{ba,cd}^{(3)}$ , we can compute explicitly the  $q$ -determinant of  $K$  in the following realizations

1. for the matrix  $K = XZ^\dagger$  (and hence for the  $q$ -twistor  $K = XX^\dagger$ )

$$\begin{aligned}
(det_q K)P_{-ij,kl} &= P_{-ij,ab}K_{ac}\hat{R}_{cb,mn}^{(3)}K_{mp}P_{-pn,kl} \\
&\propto \epsilon_{ij}^q\epsilon_{ab}^qX_aZ_c^\dagger R_{bc,mn}^{(3)}X_mZ_p^\dagger\epsilon_{pn}^q\epsilon_{kl}^q \\
&= \epsilon_{ij}^q\epsilon_{ab}^qX_aX_bZ_n^\dagger Z_p^\dagger\epsilon_{pn}^q\epsilon_{kl}^q \\
&= \epsilon_{ij}^q(X^t\epsilon^q X)(Z^t\epsilon^q Z)^\dagger\epsilon_{kl}^q = 0
\end{aligned} \tag{32}$$

since  $(X^t\epsilon^q X) = 0 = (Z^t\epsilon^q Z)$ . This reflects the well-known fact in non deformed twistor theory that twistors constructed out of two spinors determine null length vectors;

2. for the  $q$ -twistor  $K = XZ^\dagger + ZX^\dagger$ , a similar calculus to the previous one gives

$$\begin{aligned}
(det_q K)P_{-ij,kl} &\propto \epsilon_{ij}^q\epsilon_{ab}^q(X_aZ_c^\dagger + Z_aX_c^\dagger)R_{bc,mn}^{(3)}(X_mZ_p^\dagger + Z_mX_p^\dagger)\epsilon_{pn}^q\epsilon_{kl}^q \\
&= \epsilon_{ij}^q[(X^t\epsilon^q Z)(X^t\epsilon^q Z)^\dagger + (Z^t\epsilon^q X)(Z^t\epsilon^q X)^\dagger]\epsilon_{kl}^q \neq 0
\end{aligned} \tag{33}$$

then, to get twistors with non-null  $q$ -determinant we need four spinors in the definition of  $K$  (notice that  $X, X^\dagger, Z$  and  $Z^\dagger$  are algebraically independent objects). If the scalar products  $(X^t\epsilon^q Z)$  and  $(Z^t\epsilon^q X)$  are central elements in the algebra generated by  $X, Z, X^\dagger$  and  $Z^\dagger$  the  $q$ -determinant of  $K$  is also central.

### 3 An application: $q$ -Minkowski space

The classical construction of a Minkowski vector uses two (dotted and undotted) spinors,

$$K_{\alpha\dot{\beta}} = \xi_\alpha\xi_{\dot{\beta}} = (\sigma_\mu x^\mu)_{\alpha\dot{\beta}} \quad \alpha, \dot{\beta} = 1, 2 \quad , \tag{34}$$

and  $K'_{\alpha\dot{\beta}} = A_\alpha{}^\gamma K_{\gamma\dot{\delta}}(\tilde{A}^{-1})^{\dot{\delta}}{}_{\dot{\beta}}$ , where  $A$  and  $\tilde{A} = (A^{-1})^\dagger$  are the two fundamental representations of  $SL(2, C)$ . A  $q$ -deformation of the Lorentz group may be obtained [10]-[12] by replacing  $A$  and  $\tilde{A}$  by two copies  $T$  and  $\tilde{T}$  of  $SL_q(2, C)$ . Applying the pattern described above we now have two pairs of  $q$ -spinors

$$X \rightarrow X' = TX \quad Z \rightarrow Z' = TZ \quad , \tag{35}$$

$$X^\dagger \rightarrow X'^\dagger = X^\dagger\tilde{T}^{-1} \quad Z^\dagger \rightarrow Z'^\dagger = Z^\dagger\tilde{T}^{-1} \quad , \tag{36}$$

obviously, the reality condition  $T^\dagger = \tilde{T}^{-1}$  must be considered to have that  $(X')^\dagger = X'^\dagger$ , from which we may construct the following hermitian objects ( $q$ -twistors)

$$K = XX^\dagger \quad \text{or} \quad K = XZ^\dagger + ZX^\dagger \quad , \tag{37}$$



and find their transformation properties. When the reality condition  $T^\dagger = \tilde{T}^{-1}$  is imposed, the coaction

$$K' = TK\tilde{T}^{-1} = TKT^\dagger \quad (38)$$

preserves the hermiticity property of  $K$ . Since, by assumption,  $T$  and  $\tilde{T}$  are  $SL_q(2, C)$  matrices, *i.e.*,

$$R_{12}T_1T_2 = T_2T_1R_{12} \quad , \quad R_{12}\tilde{T}_1\tilde{T}_2 = \tilde{T}_2\tilde{T}_1R_{12} \quad , \quad (39)$$

the first and last equations of (15) are fulfilled if

$$\begin{aligned} R^{(1)} &= R_{12} \quad \text{or} \quad R_{21}^{-1} \quad (\kappa_1 = q \quad \text{or} \quad q^{-1}) \quad , \\ R^{(4)} &= R_{21} \quad \text{or} \quad R_{12}^{-1} \quad (\kappa_2 = q^{-1} \quad \text{or} \quad q) \quad . \end{aligned} \quad (40)$$

Then, the basic relations which define the non-commutative algebra generated by the entries of  $K$  are given by eq. (21), which gives the following possibilities

$$R_{12}K_1R^{(2)}K_2 = K_2R^{(3)}K_1R_{21} \quad . \quad (41)$$

$$R_{12}K_1R^{(2)}K_2 = q^2K_2R^{(3)}K_1R_{12}^{-1} \quad . \quad (42)$$

As we have already discussed the second possibility (42) is not valid for the twistor with four spinors (second expression in (37)) since it does not correspond to  $R^{(1)} = R^{(4)t}$ . However, it is easy to check that the algebra generated by the entries of  $K$  satisfying eq. (42) coincides with the algebra determined by (41) with the additional condition  $\det_q K = 0$ . To see it, the following consequences of the eigenvalue decomposition of  $R$  ( $\hat{R} \equiv \mathcal{P}R = qP_+ - q^{-1}P_-$ ) are useful

$$P_- \hat{R} = \hat{R} P_- = -q^{-1}P_- \quad , \quad P_- \hat{R}^{-1} = \hat{R}^{-1} P_- = -qP_- \quad , \quad (43)$$

$$q^2 \hat{R}^{-1} = \hat{R} - (q^3 - q^{-1})P_- \quad . \quad (44)$$

Multiplying now eq. (42) by  $P_- \mathcal{P}$  from the left and by  $\mathcal{P} P_-$  from the right and using (43) we get

$$-q^{-1}P_- K_1 R^{(2)} K_2 \mathcal{P} P_- = -q^3 P_- \mathcal{P} K_2 R^{(3)} K_1 P_- \quad (45)$$

as  $\hat{R}^{(i)} = \mathcal{P} R^{(i)}$ ,  $R^{(2)} = \mathcal{P} R^{(3)} \mathcal{P}$  and  $K_1 = \mathcal{P} K_2 \mathcal{P}$

$$(q^3 - q^{-1})P_- K_1 \hat{R}^{(3)} K_1 P_- = 0 \quad . \quad (46)$$

Thus, (if  $q^4 \neq 1$ ) we obtain that  $\det_q K P_- = P_- K_1 \hat{R}^{(3)} K_1 P_- = 0$ .

Now, using (44) the RE (42) can be expressed in the following way

$$R_{12}K_1R^{(2)}K_2 = K_2R^{(3)}K_1R_{21} - (q^3 - q^{-1})P_- K_2 R^{(3)} K_1 P_- \mathcal{P} \quad (47)$$

and using the definition of the  $q$ -determinant

$$R_{12}K_1R^{(2)}K_2 = K_2R^{(3)}K_1R_{21} - (q^3 - q^{-1})\mathcal{P}(\det_q K)P_- \mathcal{P} \quad (48)$$

as  $\det_q K = 0$ , we just obtained eq. (44); thus, as the algebra is the same eq. (42) may be discarded.

The matrices  $R^{(2)}, R^{(3)}$  are not determined, and characterize the mixed commutation relations between quantum group elements and conjugated elements in (16), (23) and

$$\tilde{T}_1^{-1} R^{(2)} T_2 = T_2 R^{(2)} \tilde{T}_1^{-1} \quad , \quad T_1 R^{(3)} \tilde{T}_2^{-1} = \tilde{T}_2^{-1} R^{(3)} T_1 . \quad (49)$$

Two particularly special cases arise

**a) Commuting case:** if the two quantum group copies are independent, the quantum matrices commute

$$T_1 \tilde{T}_2 = \tilde{T}_2 T_1 \quad , \quad (50)$$

here,  $R^{(2)} = R^{(3)} = I$ , and then, eq. (41) gives the reflection equation, which is equivalent to the ‘ $RTT$ ’ relation (18) (see below)

$$R_{12} K_1 K_2 = K_2 K_1 R_{21} \quad (51)$$

Eq. (42), in this particular case, produces the RE

$$R_{12} K_1 K_2 = q^2 K_2 K_1 R_{12}^{-1} \quad , \quad (52)$$

however, as we have just shown, this possibility leads to the same commutation relations (51) for the entries of  $K$  plus the additional condition  $\det_q K = 0$  [8]. The  $q$ -Minkowski algebra (51) is isomorphic to the quantum group algebra  $GL_q(2)$ , by [8]

$$T = K \sigma^1 \quad , \quad R_{12} T_1 T_2 = T_2 T_1 R_{12} \quad , \quad (53)$$

where  $\sigma^1$  is the usual Pauli matrix. Then, it is not possible to define a linear central element in the algebra generated by the entries  $(\alpha, \beta, \gamma, \delta)$  of matrix  $K$ , and the quadratic one is given by the  $q$ -determinant (28) with  $\hat{R}^{(3)} = \mathcal{P} R^{(3)} = \mathcal{P}$

$$\det_q K P_- = P_- K_1 \mathcal{P} K_1 P_- = (-q^{-1})(\alpha\delta - q\gamma\beta) P_- \quad . \quad (54)$$

**b) Non-commuting case:** now, assuming the non-trivial commutation relations between the two copies of  $SL_q(2, C)$

$$R_{12} T_1 \tilde{T}_2 = \tilde{T}_2 T_1 R_{12} \quad (55)$$

we see that (49) is fulfilled for  $R^{(2)} = R_{21}$ . Then, eq. (44) leads to the RE

$$R_{12} K_1 R_{21} K_2 = K_2 R_{12} K_1 R_{21} \quad (56)$$

Again, (42) produces an equation

$$R_{12} K_1 R_{21} K_2 = q^2 K_2 R_{12} K_1 R_{12}^{-1} \quad (57)$$

which leads to the same commutation relations as (56) with the restriction  $\det_q K = 0$ .

These equations [8] define the quantum Minkowski algebra of [10]-[12], in which the linear central term is identified with the time coordinate and the  $q$ -determinant, defined by (28) where  $\hat{R}^{(3)} = \hat{R}$

$$\det_q K P_- = P_- K_1 \hat{R} K_1 P_- = (-q^{-1})(\alpha\delta - q^2\beta\gamma)P_- \quad , \quad (58)$$

gives the quadratic central element which is identified with the invariant  $q$ -Minkowski length.

Having a  $q$ -vector  $X \mapsto TX$  and a  $q$ -matrix  $K \mapsto TKT^\dagger$ , it is natural to construct higher rank tensors transforming as

$$\varphi : L \longmapsto T^{\otimes n} L (T^\dagger)^{\otimes n} \quad ; \quad (59)$$

they are invariant subspaces of the  $q$ -Minkowski algebra for the coaction  $\varphi$ . The generators of the  $q$ -tensors  $L$  may be extracted from matrices of higher dimensions, *e.g.*

$$\begin{aligned} L^2 \sim K^{\otimes_q 2} &= K_1 R_{21} K_2 \quad \longmapsto T_1 T_2 K^{\otimes_q 2} T_1^\dagger T_2^\dagger \quad , \\ L^3 \sim K^{\otimes_q 3} &= K_1 R_{21} K_2 R_{31} R_{32} K_3 \quad , \end{aligned} \quad (60)$$

$$L^n \sim K^{\otimes_q n} = K_1 \prod_{j=2}^n (R_{j1} R_{j2} \dots R_{j,j-1} K_j) \quad .$$

These subspaces (as in the non-deformed theory) are reducible (for instance,  $K^{\otimes_q 2}$  has  $\det_q K$  as an invariant element). One can apply to  $T^{\otimes 2} = T_1 T_2$  the appropriate projector  $P^{(1)}$  to the spin 1 representation (and the same for  $(T^\dagger)^{\otimes 2}$ ) to get a tensor of generators transforming according  $D^{1,1}$  *irrep* of the  $q$ -Lorentz group. Quantum tensors transforming according  $D^{j,s}$  *irreps* could be constructed in the same manner. We find additional  $R$ -matrix factors in the tensor products of  $K$  (60) (cf. (28)). This construction is useful for a description of higher spin  $q$ -wave equations (see also [13]).

**Acknowledgements:** This research has been partially sponsored by a CICYT (Spain) research grant. One of us (P.P.K.) also wishes to thank the DGICYT, Spain, for financial support.

## References

- [1] V.G. Drinfel'd, in Proc. of the 1986 *Int. Congr. of Mathematicians*, MSRI Berkeley, vol. **I**, 798 (1987)
- [2] M. Jimbo, Lett. Math. Phys. **10**, 63 (1985); *ibid.* **11**, 247 (1986)
- [3] L.D. Faddeev, N. Yu. Reshetikhin and L.A. Takhtajan, Alg. i Anal. **1**, 178 (1989) (Leningrad Math. J. **1**, 193 (1990))
- [4] Yu. I. Manin, Commun. Math. Phys. **123**, 163 (1989); *Topics in non-commutative geometry*, Princeton Univ. Press. (1991)
- [5] J. Wess and B. Zumino, Nucl. Phys. (Proc. Suppl.) **18B**, 302-312 (1990)
- [6] P.P. Kulish and E.K. Sklyanin, J. Phys. **A25**, 5963 (1992)
- [7] P.P. Kulish and R. Sasaki, Progr. Theor. Phys. **89**, 741 (1993)
- [8] J.A. de Azcárraga, P.P. Kulish and F. Rodenas, Lett. Math. Phys. (1994); *Quantum groups and deformed special relativity*, FTUV-94-21/IFIC-94-19 (April 1994, hep-th/9405161)
- [9] Majid S., in *Quantum Groups*, Lect. Notes Math. **1510**, 79 (1992); J. Math. Phys. **32**, 3246 (1991); *ibid* **34**, 1176 (1993)
- [10] U. Carow-Watamura, M. Schlieker, M. Scholl and S. Watamura, Z. Phys. **C48**, 159 (1990); Int. J. Mod. Phys. **A6**, 3081 (1991)
- [11] W. Schmidke, J. Wess and B. Zumino, Z. Phys. **C52**, 471 (1991)
- [12] O. Ogievetsky, W. B. Schmidke, J. Wess and B. Zumino, Commun. Math. Phys. **150**, 495 (1992)
- [13] M. Pillin, J. Math. Phys. **35**, 2804 (1994)